AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in this application.

Listing of Claims:

1. (original) A process for the acylation of a C(10) hydroxy group of a taxane, the process comprising

treating the taxane with an acylating agent in a reaction mixture containing less than one equivalent of an amine base for each equivalent of taxane to form a C(10) acylated taxane.

- 2. (original) The process of claim 1 wherein the taxane has C(7) and C(10) hydroxy groups and the acylating agent selectively reacts with the C(10) hydroxy group.
- 3. (original) The process of claim 1 wherein the taxane reacted with the acylating agent is 10-deacetyl baccatin III.
 - 4. (currently amended) The process of claim 1 wherein the taxane has the structure:

wherein

 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;

 R_2 is keto, -OT₂, acyloxy, or together with R_1 forms a carbonate;

 R_4 is -OT₄ or acyloxy;

 R_7 is hydrogen, halogen, $-OT_7$, $-OCOZ_7$, or $-OCOOZ_7$;

 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, or $-OCOOZ_9$;

 R_{10} is hydroxy;

R₁₃ is hydroxy, protected hydroxy, keto, or

$$X_{5} \xrightarrow{C} X_{4} \xrightarrow{X_{3}} 0$$

$$H X_{2} \xrightarrow{X_{1}} 0$$

 R_{14} is hydrogen, $-OT_{14}$, acyloxy, or together with R_1 forms a carbonate;

 T_2 , T_4 , T_7 , T_9 and T_{14} are independently hydrogen or hydroxy protecting group;

 X_1 is $-OX_6$, $-SX_7$, or $-NX_8X_9$;

 X_2 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

$$X_5$$
 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, $-NX_8X_{10}$, or $-SO_2X_{11}$;

 X_6 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

 X_7 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, or sulfhydryl protecting group;

 X_8 is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

 X_9 is an amino protecting group;

X₁₀ is hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

X₁₁ is hydrocarbyl, substituted hydrocarbyl, heteroaryl, -OX₁₀, or -NX₈X₁₄; [[and]]

 X_{14} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl[[.]]; and

Z₇ is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

Z₉ is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

5. (original) The process of claim 4 wherein

 R_1 is hydroxy or together with R_{14} or R_2 forms a carbonate;

 R_2 is $-OCOZ_2$, $-OCOOZ_2$, or together with R_1 forms a carbonate;

 R_4 is $-OCOZ_4$;

R₉ is hydrogen or keto;

R₁₃ is hydroxy, protected hydroxy, or

 R_{14} is hydrogen, hydroxy, protected hydroxy, or together with R_1 forms a carbonate;

 X_1 is $-OX_6$ or $-NX_8X_9$;

X₂ is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

 X_5 is $-X_{10}$, $-OX_{10}$, or $-NX_8X_{10}$;

X₆ is a hydroxy protecting group;

X₈ is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X₉ is an amino protecting group;

 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

 Z_2 and Z_4 are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

FSUM 10302 PATENT

- 6. (original) The process of claim 1 wherein the acylating agent is selected from the group consisting of anhydrides, dicarbonates, thiodicarbonates, and isocyanates.
 - 7. (original) The process of claim 6 wherein the reaction mixture contains a Lewis acid.
- 8. (original) The process of claim 7 wherein the taxane reacted with the acylating agent is 10-deacetyl baccatin III.
 - 9. (original) The process of claim 1 wherein the reaction mixture contains a Lewis acid.
- 10. (original) The process of claim 9 wherein the taxane reacted with the acylating agent is 10-deacetyl baccatin III.
 - 11. (currently amended) The process of claim 9 wherein the taxane has the structure:

wherein

 R_1 is hydrogen, hydroxy, protected hydroxy, or together with R_{14} or R_2 forms a carbonate;

R₂ is keto, -OT₂, acyloxy, or together with R₁ forms a carbonate;

R₄ is -OT₄ or acyloxy;

 R_7 is hydrogen, halogen, $-OT_7$, $-OCOZ_7$, or $-OCOOZ_7$;

 R_9 is hydrogen, keto, $-OT_9$, $-OCOZ_9$, or $-OCOOZ_9$;

R₁₀ is hydroxy;

 R_{13} is hydroxy, protected hydroxy, keto, or

FSUM 10302 PATENT

 R_{14} is hydrogen, -OT₁₄, acyloxy, or together with R_1 forms a carbonate;

T₂, T₄, T₇, T₉ and T₁₄ are independently hydrogen or hydroxy protecting group;

 X_1 is -OX₆, -SX₇, or -NX₈X₉;

X₂ is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

 X_5 is $-X_{10}$, $-OX_{10}$, $-SX_{10}$, $-NX_8X_{10}$, or $-SO_2X_{11}$;

 X_6 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, hydroxy protecting group or a functional group which increases the water solubility of the taxane derivative;

 X_7 is hydrocarbyl, substituted hydrocarbyl, heteroaryl, or sulfhydryl protecting group;

X₈ is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

 X_9 is an amino protecting group;

 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

 X_{11} is hydrocarbyl, substituted hydrocarbyl, heteroaryl, $-OX_{10}$, or $-NX_8X_{14}$; [[and]]

 X_{14} is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl[[.]]; and

 Z_7 is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

Z₉ is hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

12. (original) The process of claim 11 wherein

R₁ is hydroxy or together with R₁₄ or R₂ forms a carbonate;

 R_2 is $-OCOZ_2$, $-OCOOZ_2$, or together with R_1 forms a carbonate;

 R_4 is $-OCOZ_4$;

R₉ is hydrogen or keto;

R₁₃ is hydroxy, protected hydroxy, or

 R_{14} is hydrogen, hydroxy, protected hydroxy, or together with R_1 forms a carbonate;

 X_1 is $-OX_6$ or $-NX_8X_9$;

X₂ is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

 X_3 and X_4 are independently hydrogen, hydrocarbyl, substituted hydrocarbyl, or heteroaryl;

 X_5 is $-X_{10}$, $-OX_{10}$, or $-NX_8X_{10}$;

 X_6 is a hydroxy protecting group;

X₈ is hydrogen, hydrocarbyl, or substituted hydrocarbyl;

X₉ is an amino protecting group;

 X_{10} is hydrocarbyl, substituted hydrocarbyl, or heteroaryl; and

 Z_2 and Z_4 are independently hydrocarbyl, substituted hydrocarbyl, or heteroaryl.

- 13. (original) The process of claim 9 wherein the Lewis acid is selected from the group consisting of the halides or triflates of the Group IB, IIB, IIIB, IVB, VB, VIIB, VIII, IIIA, IVA, lanthanide and actinide elements.
- 14. (original) The process of claim 13 wherein the Lewis acid is selected from the group consisting of zinc chloride, stannic chloride, cerium trichloride, cuprous chloride, lanthanum trichloride, dysprosium trichloride and ytterbium trichloride.

- 15. (original) The process of claim 1 wherein the C(10) acylated taxane comprises a C(7) hydroxy group and the process additionally comprises treating the C(10) acylated taxane with a silylating agent to silylate the C(7) hydroxy group.
 - 16. (original) The process of claim 15 wherein the C(10) acylated taxane is baccatin III.
- 17. (original) The process of claim 1 wherein the C(10) acylated taxane comprises a C(7) hydroxy group and the process additionally comprises treating the C(10) acylated taxane with an acylating agent to acylate the C(7) hydroxy group.
 - 18. (original) The process of claim 17 wherein the C(10) acylated taxane is baccatin III.
- 19. (original) The process of claim 1 wherein the C(10) acylated taxane comprises a C(13) hydroxy, metallic oxide, or ammonium oxide substituent and the process additionally comprises the step of esterifying the C(10) acylated taxane by treating the C(10) acylated taxane with a side chain precursor selected from the group consisting of β -lactams, oxazolines, oxazolidine carboxylic acids, oxazolidine carboxylic acid anhydrides, and isoserine derivatives.

Claims 20-45 (cancelled)